## **ACTH Stimulation for Dogs**

| Drug           | Synthetic ACTH (Cortrosyn<br>(Cosyntropin for injection))  | Synthetic ACTH (Synacthen Depot (Cosyntropin-Zinc Hydroxide))             | ACTH gel (Bexco (Corticotropin injection, USP)) |
|----------------|--|---|---|
| Formulation    | 0.25 mg/vial reconstituted with 1<br>mL sterile saline,<br>0.25 mg = 250 μg = 25 U   | 1 mg/vial, 1 mL bottle  | 40 U/mL, 5 mL bottle                            |
| Dose           | 250 μg (1 vial)/dog regardless of<br>body weight <b>or</b> (Cost-effective<br>lower dose of 5 μg/kg to a<br>maximum of 250 μg/dog) | body weight <15 kg -<br>0.25 mg/dog<br>body weight >15 kg -<br>0.5 mg/dog | 2.2 U/kg of body weight                         |
| Administration | intravenously  | intramuscularly   | intramuscularly                                 |
| Collection     | Pre and 1 hour post  | Pre and 2 hour post   | Pre and 2 hours post                            |

## **ACTH Stimulation for Cats**

| Drug           | Synthetic ACTH (Cortrosyn (Cosyntropin for injection))                             | Synthetic ACTH (Synacthen Depot (Cosyntropin-Zinc Hydroxide)) | ACTH gel (Bexco (Corticotropin injection, USP)) |
|----------------|--|---|---|
| Formulation    | 0.25 mg/vial reconstituted with<br>1 mL sterile saline,<br>0.25 mg = 250 μg = 25 U | No established protocol                                       | 40 U/mL, 5 mL bottle                            |
| Dose           | 125 μg (½ vial)/cat regardless of body weight                                      |   | 2.2 U/kg of body<br>weight                      |
| Administration | intravenously  |   | intramuscularly                                 |
| Collection     | Pre, 30 min. and 1 hour post   |   | Pre, 1 and 2 hours post                         |